

*Application
for
United States Letters Patent*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
22

**Assays For Evaluating
The Function Of RNA Helicases**

5

The invention disclosed herein was made with Government support under NIH Grant No. RO150313 from the Department of Health and Human Services. Accordingly, the U.S. Government has certain rights in this invention.

10

Background on the Invention

15

Throughout this application, various publications are referenced by author and date. Full citations for these publications may be found listed alphabetically at the end of the specification immediately preceding Sequence Listing and the claims. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art as known to those skilled therein as of the date of the invention described and claimed herein.

20

25

RNA conformation is of elemental importance in RNA-induced catalysis, as well as RNA interactions with other cellular components. Many central components of gene expression and RNA metabolism occur in large ribonucleoprotein complexes, most notably the ribosome and spliceosome. Within these large complexes, alternative conformations of specific RNAs have been demonstrated and the alteration of RNA conformation is believed to play a critical role in enabling, driving and assuring the fidelity of the catalytic reactions these complexes perform (Staley and Guthrie, 1998). Therefore, understanding the factors capable of inducing RNA conformational alterations within these complexes is likely to be central to an overall appreciation of

30
35

mechanisms by which they execute their intricate and demanding functions.

By exploring the effect of the duplex length of the substrates on the unwinding reaction under defined reaction conditions, the response of unwinding rate and amplitude to the duplex length can provide information about processivity and directionality (Fig. 1). Thus, the requirements for an experimental setup to probe and quantify processivity and directionality are to develop a substrate system where only the duplex length is variable and, using this substrate system, establish experimental conditions where rate and/or amplitude provide information about processivity and directionality.

RNA helicases of the DExH/D family play an essential role in viral replication and cellular RNA metabolism, including central functions in RNA splicing, translation and regulation of gene expression. Despite the importance of these proteins, their RNA helicase activity has not been subjected to enzymological study. Basic knowledge of cellular metabolism is therefore constrained by a limited understanding of reaction mechanism by motor proteins in the RNA helicase family. To address this problem, mechanistic studies have been initiated on two viral DExH/D proteins which are part of the RNA helicase family: NPH-II from Vaccinia and NS3-4A from Hepatitis C Virus (HCV). The NPH-II protein is shown to be a processive, directional RNA helicase with specific roles for both the binding and hydrolysis of ATP. Having established qualitative features of NPH-II activity, the use of direct and stopped-flow kinetic methods to determine the quantitative kinetic parameters such as

translocation rates, reaction step size, processivity, helicase binding, ATP binding and hydrolytic rate constants that describe the framework for catalytic activity of this prototypical RNA helicase. All aspects of cellular RNA metabolism and processing involve DExH/D proteins, which are a family of enzymes that unwind or manipulate RNA in an ATP-dependent fashion (de la Cruz, et al., 1999). DExH/D proteins are also essential for the replication of many viruses, and therefore provide targets for the development of therapeutics (Radare and Haenni, 1999). All DExH/D proteins characterized to date hydrolyse nucleoside triphosphates and, in most cases, this activity is stimulated by the addition of RNA or DNA. Several members of the family unwind RNA duplexes in an NTP dependent fashion *in vitro* (de la Cruz, et al., 1999 and Wagner, et al., 1998); therefore it has been proposed that DExH/D proteins couple NTP hydrolysis to RNA conformational change in complex macromolecular assemblies (Stanley and Guthrie, 1998). Despite the central role of DExH/D proteins, their mechanism of RNA helicase activity remains unknown. It is shown that the DExH protein NPH-II unwinds RNA duplexes in a processive, unidirectional fashion with a step size of roughly one-half helix turn and that there is a quantitative connection between ATP and helicase processivity, thereby providing direct evidence that DExH/D proteins can function as molecular motors on RNA.

Summary of the Invention

The present invention provides a method for detecting the release of a single-stranded RNA from an RNA duplex which comprises: (a) admixing an RNA helicase with the RNA duplex under conditions permitting the RNA helicase to unwind the RNA duplex and release single-stranded RNA, wherein the RNA duplex comprises a first RNA having a first label attached thereto and a second RNA, such first label being capable of producing a luminescent energy pattern when the first RNA is present in the RNA duplex which differs from the luminescent energy pattern produced when the first RNA is not present in the RNA duplex, (b) subjecting the admixture to conditions which permit (i) the RNA helicase to unwind the RNA duplex and release single-stranded RNA, and (ii) the first label to produce luminescent energy; and (c) detecting a change in the luminescent energy pattern produced by the first label so as to thereby detect release of single-stranded RNA from the RNA duplex.

Brief

Brief Description of the Figures

Figure 1

Kinetic approach to probe processivity and directionality of the unwinding reaction. Unwinding of a dsRNA is the reverse reaction of the association of two complementary ssRNAs. Thus, unwinding occurs only when so many basepairs are disrupted that no nucleation locus can form. Practically, that means that unwinding is only observed, when less than 4 or 3 basepairs remain in the duplex. This emphasizes that by monitoring a duplex unwinding reaction one actually monitors the very last event of the strand separation process.

A processive reaction occurs in distinct consecutive steps, therefore, there are reaction intermediates (I) until the reaction product (P) is formed. For a non-processive reaction (for instance if the unwinding occurs in one power stroke) there are no intermediates. Thus, the experimental rationale for probing processivity is to determine whether there are reaction intermediates or not. If the reaction proceeds in distinct consecutive translocation steps, the translocation is rate limiting for the unwinding reaction, and the rate constants (k_{tr}) of these translocation steps are of similar or equal size, intermediates (I) would accumulate. This would be reflected in a lag phase in the unwinding timecourse of the product (P) formation. With increasing duplex length more reaction intermediates would appear and timecourses would display a greater lag phase with increasing duplex length. The number of intermediates ($I_{1...n}$) is indicative for the number of kinetic steps that the helicase takes to unwind a duplex of given length. More intermediates cause also more dissociation "events", provided that these intermediates are susceptible to dissociation. This is

determined by the ratio of the translocation rate constant (k_{tr}) to the respective dissociation rate constant (k_{di}) (Ali and Lohmann, 1997). If a measurable fraction of protein dissociates during each unwinding step, the reaction amplitude should decrease with increasing duplex length. This should also occur when the translocation is not rate-limiting for the unwinding reaction. However, in that case the duplex length should not affect the reaction rate. If the reaction is not processive, there should be under no conditions a sensitivity of either reaction rate or amplitude to the length of the duplexes.

Probing the processivity is the prerequisite for testing the directionality of the unwinding, since only a processive reaction can be directional. The directionality can be directly tested by monitoring at least one reaction intermediate and the reaction product at the same time under conditions where the translocation limits the unwinding reaction. The experimental conditions for probing processivity and directionality have to be chosen in a way, that all RNA is complexed with protein at the reaction start and that all dissociation of the protein off the RNA substrate is irreversible.

Figure 2

Unwinding of various RNA substrates by NPH-II. Unwinding reactions were performed in a reaction buffer comprising 30 μ L volume (40mM Tris.Cl, pH 8.0, 2mM DTT, 20mM NaCl, 3mM $MgCl_2$, 3mM ATP, 16nM NPH-II, 3nM RNA substrate) at 23 C. Protein, RNA substrate, and $MgCl_2$ were incubated in reaction buffer prior to the reaction for 7 min. The reaction was started by addition of ATP. Aliquots were

withdrawn at appropriate times and added to two volumes quenching buffer (25mM EDTA, 0.4% SDS, 10% glycerol, 0.05% BPB, 0.05% XCB). This mixture was immediately cooled on ice. After the last aliquot was taken, the samples were applied to 15% native PAGE.

Figure 3

Effect of trap RNA on unwinding reactions. (A) Since reactions are performed with preannealing of protein and RNA prior to the reaction, the protein-RNA complex [ES] is already formed at the reaction start. As protein dissociates from the RNA substrate during the unwinding reaction, re-binding of the protein and multiple reaction cycles can occur. Binding ("trapping") free protein during the course of unwinding prevents re-binding during the reaction. Complete "trapping" of all free protein is ensured by using a large excess of RNA (trap RNA) as compared to the duplex substrate. (B) Effect of trap addition on reactions with LS83 and LS36. Reaction conditions were as described in Fig. 2, except that the ATP was pre-mixed with the trap RNA (39-mer RNA oligo) and the reaction was started with this ATP/trap mixture. The trap concentration in the reaction was 600nM which is an 200 fold excess over the substrate. Higher concentrations of trap did not change the observed timecourses. The substrate was bound to 100% of the protein prior to the reaction as verified by gel shift analysis of the RNA protein complexes (not shown).

Figure 4

Monitoring duplex unwinding by fluorescence energy transfer. (A) Both strands of the duplex are labeled with a fluorescence transfer donor-acceptor-pair of fluorescent dyes. The top strand is labeled with

fluorescein (donor) at the 3' -end and the bottom strand is labeled with rhodamin (acceptor) at the 5' -end. The attached fluorescein is excited at 492nm. Due to the proximity of the acceptor label in the duplex, the emission of the fluorescein label is quenched. Upon unwinding, both labels depart from each other, the quenching of the fluorescein label no longer occurs and an increase in the emission of the fluorescein fluorescence can be detected. (B) The unwinding reactions were performed as described in Fig. 3, but in a volume of 600 μ L in a fluorescence cuvette. The fluorescence intensity was measured at defined intervals (small points). The resulting curve was corrected for the reaction amplitude derived by gel shift measurements of identical reactions and both, the derived fluorescence curve and the measurements from gel shift experiments (large points) are overlaid in the plots. Reactions were performed with US18 (18bp duplex region) and US36 (36bp duplex region).

Figure 5

Effect of ATP concentration on the unwinding reaction. Unwinding reactions with LS36 were performed as described in Fig. 3 (pre-annealing, trap addition). The plot shows the final amplitude of the reaction versus the respective ATP concentration. Amplitude values were determined out of 8 aliquotes taken from the reaction.

Figure 6

Effect of Mg^{2+} substitution on the unwinding reaction. Timecourses with US36 were measured as described in Fig. 3 (trap RNA, pre-annealing), except that 3nM Mg^{2+} (A) was substituted with 3mM Mn^{2+} (B) and 3mM Co^{2+} (C). The left panels show the gel printouts of the reactions that are

plotted on the right panels as fraction unwound substrate versus time.

Figure 7

5 Effect of the duplex length on unwinding reactions in the presence of Co^{2+} . Unwinding reactions were performed with US36 (36bp duplex region) and US18 (18bp duplex region) under the conditions described in Fig. 6. The solid lines represent fits for a consecutive 2 step reaction with equal rate constants (Ali and Lohmann, 1997) for the 18bp duplex and for a consecutive 4 step reaction with equal rate constants for the 36bp duplex.

Figure 8

15 Probing the directionality of the unwinding reaction by using multi-piece substrates. (A) Multi-piece substrates were generated by site directed processing of the top strand RNA with an engineered DNAzyme (Santoro and Joyce, 1997) and subsequent hybridization of the two "pieces" with the bottom strand. (B) Unwinding reactions of MPS 36/75 (36nt oligo binds next to the single stranded overhang and 75nt oligo binds adjacent to this strand) and MPS 83/28 (83nt oligo binds next to the single stranded overhang and 28nt oligo binds adjacent to this strand) in the presence of Mg^{2+} were carried out as described in Fig. 3. The fraction of the displaced pieces was calculated according to $\text{frac}[N]_t = [N_t / (N_t + M_t + S_t)] * [(N_\infty + M_\infty + S_\infty) / N_\infty]$, where $\text{frac}[N]_t$ is the fraction of the respective "piece" at the time t ; N_t is the intensity of the band of the "piece" N at the time t ; M_t is the intensity of the band of the "piece" M at the time t ; S_t is the intensity of the not unwound substrate band at the time t ; N_∞ is the intensity of the band of

the "piece" N after heat denaturation; M_{∞} is the intensity of the band of the "piece" M after heat denaturation and S_{∞} is the intensity of the not unwound substrate band after heat denaturation. S_{∞} is, however, usually zero. The solid lines represent a fit of the timecourse to a rate law describing a sum of two first order reactions. (C) Reactions of MPS 36/75 and MPS 83/28 in the presence of Co^{2+} were performed as described in Fig. 6. The fraction of the displaced pieces were calculated as described above. The solid lines represent fits of the timecourses to rate laws describing a consecutive 4 step reaction for T36, a 12 step reaction for T75 and T28 and a 9 step reaction for T 83, using equal rate constants for the steps, respectively.

Figure 9

Kinetics of duplex unwinding. Reactions in the presence (open circles) and absence (filled circles) of trap RNA with a 36-bp substrate (a) and a 83-bp substrate (b). The 83-bp substrate is an extension of the 36-bp substrate. The fraction of unwound substrate (indicated on the y axis) was fitted to the integrated first-order rate law using Kaleidagraph software (Abelbeck). Rate constants (k) and reaction amplitudes (A) in the absence of trap RNA: $k = 3.5 \pm 0.1 \text{ min}^{-1}$, $A=0.96$, 36-bp substrate; $k = 3.4 \pm 0.1 \text{ min}^{-1}$, $A=0.97$, 83-bp substrate. In the presence of trap RNA: $k = 3.6 \pm 0.2 \text{ min}^{-1}$, $A=0.77$, 36-bp substrate; $k = 3.6 \pm 0.3 \text{ min}^{-1}$, $A=0.52$, 83-bp substrate. Variances are standard deviations from the fit.

Figure 10

Estimation of the unwinding step size. Time courses of substrates containing duplex regions of 12 (open squares), 18 (filled squares), 24 (open circles) and 36

(filled circles) bp unwound by NPH-II in the presence of 4mM CoCl_2 . Substrates contain a 35-U single-strand overhang 3' to the duplex region. Step size and the unwinding (translocation) rate constant were calculated were calculated from at least two independent time courses for each duplex which are overlaid on the plots. Solid lines are best fits to the kinetic model resulting in an unwinding step size of 6 bp and an average unwinding rate constant of $k_g = 13.4 \pm 0.8 \text{ min}^{-1}$.

Figure 11

Probing helicase directionality with multipiece substrates (MPS) **a**, Design of the MPS **b**. Unwinding of MPS in 4mM MgCl_2 and trap RNA. Unwinding rate constants were similar for all pieces ($k=3.3 \pm 0.5 \text{ min}^{-1}$); reaction amplitudes (A) were MPS 36/75, T36 A=0.74; T75, A = 0.32; MPS 83/28, T83, A=0.43; T28, A=0.27. **c**. Unwinding of multipiece substrates in 4 mM CoCl_2 and trap RNA. Kinetic data were analysed as described in Fig. 2, which resulted in average values of $k_g = 12.4 \pm 2.1 \text{ min}^{-1}$ and a step size of 6pb, consistent with the values obtained for oligomers shown in Fig. 2.

Figure 12

Processivity of NPH-II **a**, ATP dependence of processivity in 4 mM MgCl_2 (filled circles) and 4mM CoCl_2 (open circles). The solid line is a hyperbolic fit: $P = P_{\infty}[\text{ATP}] / (Z + [\text{ATP}])$, where $P_{\infty} = k_{U(S)} / (k_{U(S)} + k_8)$ reflects the processivity at ATP saturation; Z is the ATP concentration where $P = P_{\infty}/2$. In 4mM CoCl_2 , $P_{\infty} = 1.00 \pm 0.03$ and $Z = 0.50 \pm 0.06 \text{ mM}$. In 4mM MgCl_2 , $P_{\infty} = 0.99 \pm 0.01$ and $Z = 0.14 \pm 0.01 \text{ mM}$. Processivities were determined from the reaction amplitudes of four duplexes

(12, 18, 24 and 36 bp, Fig. 2) at the ATP concentrations indicated. Amplitudes were obtained from at least three independent experiments. Processivity was calculated by fitting plots of amplitude at constant ATP concentration versus duplex length to Equation 2 using a step size of $m=6$ (Fig. 2). **b,c**, Dependence of reaction amplitude on ATP concentration for substrates with duplex regions of 12 (filled circles), 18 (open circles), 24 (filled squares) and 36 (open squares) basepairs in 4 mM $MgCl_2$ (**b**) and 4mM $CoCl_2$ (**c**). Each reaction amplitude was determined in triplicate and the deviation from these independent measurements is indicated by the error bars. The solid lines are fits to equation 1, where k'_d was assumed to equal K_m from multiple turnover ATPase measurements (1.1 mM in Mg^{2+} and 3.5 mM in Co^{2+} (Shuman, 1993; Gross and Shuman 1996), the step size was 6bp (Fig. 2), and $k_p/k_{U[S]}$ was allowed to float. In both Mg^{2+} and Co^{2+} $k_p/k_{U[S]} = 0.12$.

Figure 13

The kinetic scheme for RNA helicase activity by NPH-II. Unwinding initiation and two unwinding/translocation steps are depicted. ES_1 and ES_2 represent the enzyme substrate complex in non-ATP-bound states, while $[ES_1-ATP]$ and $[ES_2-ATP]$ represent ATP-bound states. Note that 'ATP-bound' indicates that ATP is bound but not hydrolysed, whereas 'non-ATP-bound' indicates that no nucleotide is bound or that ADP and/or inorganic phosphate are bound. The equilibrium between ATP-bound and non-ATP-bound states is fast compared with translocation with the unwinding rate constant $k_{U[S]}.k_p$ (Shuman, 1993; Gross and Shuman, 1996) is the rate constant for enzyme dissociation from substrate in the ATP-free state, dissociation in the ATP-bound state is

5

[illegible]

Detailed Description of the Invention

The present invention provides an a method for detecting the release of a single-stranded RNA from an RNA duplex which comprises (a) admixing an RNA helicase with the RNA duplex under conditions permitting the RNA helicase to unwind the RNA duplex and release single-stranded RNA, wherein the RNA duplex comprises a first RNA having a first label attached thereto and a second RNA, such first label being capable of producing a luminescent energy pattern when the first RNA is present in the RNA duplex which differs from the luminescent energy pattern produced when the first RNA is not present in the RNA duplex, (b) subjecting the admixture to conditions which permit (i) the RNA helicase to unwind the RNA duplex and release single-stranded RNA, and (ii) the first label to produce luminescent energy, and (c) detecting a change in the luminescent energy pattern produced by the first label so as to thereby detect release of single-stranded RNA from the RNA duplex.

In accordance with the present invention conditions which permit the RNA helicase to unwind the RNA duplex and release single-stranded RNA preferably include, but are not limited to the presence of ATP and a divalent cation, which preferably may be Mg^{2+} , Mn^{2+} or Co^{2+} .

The foregoing method may be carried out in numerous ways which will be readily understood by those skilled in the art. In a presently preferred embodiment of the invention the method utilizes a second label attached to the second RNA such that the luminescent energy produced by the first label interacts with the second label to produce a characteristic pattern when the first and second RNA are present in an RNA duplex, but interacts

5 differently or not at all when the RNA duplex unwinds.
In one embodiment of the invention the first label is
covalently attached at the 5' end of the first RNA and
the second label, if any, is attached at the 3' end of
the second RNA.

10 In accordance with the foregoing, embodiment the first
and second label may comprise different fluorophors
having the characteristic that the second label absorbs
luminescent energy released from the first fluorophor
when it is induced to emit luminescent energy by exposure
to a particular type of radiation such as ultraviolet
light of a defined wavelength. In one embodiment the
15 first label is fluorescein isothiocyanate and the second
label is rhodamine isothiocyanate.

20 As used herein luminescent energy includes but is not
limited to fluorescence, phosphorescence, and
chemiluminescence.

25 In the practice of the methods of the invention the
compound capable of inhibiting unwinding activity of an
RNA helicase may be a chemically synthesized substrate
that preferentially binds over a endogenous substrate to
the RNA helicase. In one embodiment of the invention the
30 compound inhibits the accessibility of the ATP to the
binding site on the protein thereby enabling the helicase
to dissociate from the RNA at appropriate times. In
another embodiment of the invention the compound induces
a rate-limiting step by blocking and deblocking
initiation of the RNA unwinding by the RNA helicase.

35 In the practice of the methods of the invention the
compound capable of inhibiting the unwinding activity of
an RNA helicase is present at a concentration capable of

inhibiting the unwinding activity of an RNA helicase. Accordingly, the effective amount will vary with the helicase used.

5 This invention also provides a method of measuring the rate of release of a single-stranded RNA from an RNA duplex which comprises detecting whether the single-stranded RNA is released from the RNA duplex at
10 predetermined time intervals according to the method and determining therefrom the rate of release of the single-stranded RNA from the RNA duplex.

15 Furthermore, this invention provides a method of determining whether a compound is capable of modulating the release of a single-stranded RNA from an RNA duplex by an RNA helicase which comprises detecting the release of the single-stranded RNA from the RNA duplex according to the method wherein the compound is added to the mixture of RNA duplex and RNA helicase

20 Still further, this invention also provides effective amounts of compounds capable of inhibiting unwinding activity of an RNA helicase together with suitable diluents, preservatives, solubilizers, emulsifiers, 25 adjuvants and/or carriers useful in treatment of viral RNA helicase or a viral Hepatitis C. Such compositions are liquids or lyophilized or otherwise dried formulations and include diluents of various buffer content (e.g., Tris-HCl., acetate, phosphate), pH and
30 ionic strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimerosal, benzyl
35

alcohol, parabens), bulking substances or tonicity modifiers (e.g., lactose, mannitol), covalent attachment of polymers such as polyethylene glycol to the compound, complexation with metal ions, or incorporation of the compound into or onto particulate preparations of polymeric compounds such as polylactic acid, polglycolic acid, hydrogels, etc, or onto liposomes, micro emulsions, micelles, unilamellar or multi lamellar vesicles, erythrocyte ghosts, or spheroplasts. Such compositions will influence the physical state, solubility, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the compound. The choice of compound will depend on the physical and chemical properties of the compound capable of alleviating the symptoms of a viral RNA infection.

This invention is illustrated in the Experimental Details section which follows. These sections are set forth to aid in an understanding of the invention but are not intended to, and should not be construed to, limit in any way the invention as set forth in the claims which follow thereafter.

Experimental Details

The DExH helicase NPH-II from vaccinia virus (Shuman, 1992) was subjected to quantitative kinetic analysis that varied in composition and length. Kinetics of unwinding were monitored using gel-shift electrophoresis and fluorescence energy transfer methodologies. The RNA substrates contained a single-strand 3' overhang, which is required for NPH-II unwinding activity (Shuman, 1993). The single-strand 3' overhang was identical in length and sequence for every substrate (Fig. 9). Unwinding reactions were preformed under single-turn-over conditions with respect to the RNA substrate. A large

excess of nonspecific trap RNA (de la Cruz, 1999) was added to prevent helicase from reassociating with duplex once it falls off during the course of reaction.

5 In the absence of trap RNA, the unwinding rate and reaction amplitude (defined as the final fraction of unwound RNA) were both insensitive to duplex length (Fig. 9). Each reaction was first-order with a rate constant of $3.5 \pm 0.2 \text{ min}^{-1}$ (Fig. 9). In the presence of trap RNA, 10 the rates remained independent of duplex length however, the reaction amplitude decreased with increasing duplex length (Fig. 9). These observations provide three mechanistic insights. First, the fact that long duplexes can be unwound at all in the presence of trap RNA 15 provides strong qualitative evidence that the helicase is a processive enzyme consistent with previous multiple-turnover studies. Second, the dependence of amplitude on duplex length indicates that more protein dissociates from an RNA substrate when the duplex is longer indicating that more protein dissociates from an RNA 20 substrate when the duplex is longer, indicating that more unwinding steps may be required for a long duplex, and there is a specific degree of processivity for the NPH-II enzyme (Fig. 9). Last, the similar rates observed for 25 the unwinding of different duplex lengths indicate that individual unwinding steps do not limit the reaction rate under these conditions. Rate-limiting events during translocation would result in a length-dependent lag phase in each time course.

30 To evaluate helicase step size and to provide an additional proof of processivity, it was necessary to monitor individual unwinding steps. Conditions were sought under which helicase translocation becomes rate 35 limiting. Variation in temperature and pH did not cause

the rates to become increasingly sensitive to duplex length (E. Jankowsky et al., unpublished data); however, marked effects were observed when Mg^{2+} , which is the cofactor for ATP hydrolysis, was replaced by Co^{2+} . The unwinding reactions displayed pronounced lag phases which increased incrementally with the duplex length (Fig. 10). The data were fit numerically to a kinetic model that relates the translocation rate constant to step size and duplex length (Material and Methods). From this analysis, the unwinding step size of NPH-II was estimated to be six base pairs (bp) and the unwinding rate constant (for each translocative step in Co^{2+} at an ATP concentration of 3.5 mM), k_u , was $13.4 \pm 0.8 \text{ min}^{-1}$. Using the step size of 6 bp, it is possible to calculate a lower limit for the translocation rate constant in Mg^{2+} . In these conditions, k_u was greater than 350 min^{-1} , indicating that translocation in Mg^{2+} is at least 25-fold faster in Co^{2+} . The slower unwinding rate in Co^{2+} is not attributable to inhibition of ATPase activity, as kinetic parameters for hydrolysis of ATP are similar in Co^{2+} and Mg^{2+} (E. Jankowsky et al., unpublished data).

The existence of a defined step size indicates regularity in the translocation process; that is, unwinding is not random with respect to individual translocation rates and the number of bp displaced in one unwinding step. The step size obtained herein corresponds to about half of a helical turn and is very similar to the unwinding step size of the UvrD DNA helicase of 4 - 5 bp (Ali and Lohmann, 1997). It is important to note that the unwinding step size is an average value that may not necessarily reflect actual microscopic translocation steps of a helicase (Ali and Lohmann, 1997).

Although the above results suggest that RNA unwinding occurs in consecutive steps, the assays that we used are designed to monitor only the very last event in strand displacement (one does not see a signal until the duplex is completely unwound). Therefore, it is impossible to distinguish between directional translocation and translocation that occurs through random facilitated diffusion (Shimamoto, 1999; Kelemen and Raines, 1999). To determine which of these mechanisms applies, detection of an unwinding intermediate was needed. This was accomplished by using multipiece substrates (MPS), consisting of a continuous bottom strand that is annealed to two adjacent pieces of top-strand RNA (Fig. 3a). The top-strand pieces were generated by site-specific cleavage of a single continuous piece of top-strand RNA (Fig. 11a). The top-strand (first RNA molecule) pieces were generated by site-specific cleavage of a single continuous piece of top-strand RNA using synthetic DNAzyme endonucleases (Santoro and Joyce, 1996). The two resultant pieces had different lengths and could therefore be distinguished by their respective electrophoretic mobilities (Fig. 11b, 11c). To prevent the results from being biased by the relative length of the top-strand pieces we designed two different sets of MPS (Fig. 11a). All MBPS unwinding reactions were carried out under single-turnover conditions with respect to the RNA substrate, and in the presence of trap RNA.

NPH-II unwound both sets of MPS (Fig. 11b), indicating that the helicase can proceed through nicks in the top strand of the substrate. In the presence of Mg^{2+} , all pieces were displaced with apparent first-order kinetics with similar rate constants, $k = 3.3 \pm 0.5 \text{ min}^{-1}$ (Fig. 11b). For both sets of MPS, the piece that was bound farthest from the overhang was displaced with a lower

amplitude than the piece located closer to the single-strand 3' overhang (Fig. 11b). This implies that fewer unwinding steps were required for the displacement of the piece closer to the overhang. The fact that these effects depend exclusively on the location of the top strands relative to the single-strand 3' overhang (rather than their length) strongly suggests that the strand displacement progresses in a distinct 3'-to-5' direction with respect to the bottom strand.

The similar unwinding rates for both pieces of top-strand RNA indicate that neither a late step in the reaction, such as the final strand separation, nor translocation itself, limits the rate of the overall process in Mg^{2+} . A late rate-limiting step would result in a slower rate for unwinding of the piece farthest from the overhang because two strand separations are required for unwinding of the second piece. A rate-limiting translocation would cause a lag phase in the time courses as observed in the presence of Co^{2+} (Fig. 10). Thus in the presence of Mg^{2+} , an initiation step at the very beginning of the unwinding process limits the rate of the overall reaction.

By contrast, reactions conducted in the presence of Co^{2+} displayed a pronounced lag phase attributable to rate-limiting translocation steps (Fig. 11c). Notably, the displacement of the two pieces occurred with different lag phases. The lag was largest for the piece farthest from the overhang, reflecting an increase in the number of unwinding steps that are required as the helicase moves away from the single-strand 3' overhang. The effects were exclusively dependent on the location of the pieces relative to the overhang and not on their length indicating a distinct 3'-to-5' directionality in the

progression of unwinding with respect to the bottom strand.

5 Having established that the helicase unwinds RNA
 substrates in a processive directional fashion with a
 distinct unwinding step size it was of interest to
 correlate this process with the role of ATP. The effect
 of ATP on enzyme processivity (P) was of particular
 interest (Fig. 12). Processivity reflects the
 10 probability that the helicase will perform the next
 unwinding step rather than dissociate from the substrate
 (Lohmann and Bjoernson, 1996). Quantitation of
 processivity was carried out by determining the reaction
 amplitude in the presence of trap RNA, such that all
 15 dissociation of protein from the RNA substrate during
 unwinding was effectively irreversible. Processivity
 decreased with decreasing ATP concentration (Fig. 12a),
 which indicates that the helicase dissociates more
 readily from RNA in non-ATP-bound states than in the ATP-
 20 bound state, consistent with previous studies of RNA
 binding by NPH-II (Shuman, 1992). In addition, the
 processivity at ATP saturation approaches $P = 1$,
 regardless of whether Co^{2+} or Mg^{2+} are the cofactors for
 ATP hydrolysis (Fig. 12a). This indicates that NPH-II in
 25 the ATP-bound state does not significantly dissociate
 from the RNA substrate, and that dissociation of protein
 from RNA occurs predominantly in non-ATP-bound states.
 This is highly instructive because it shows that ATP
 binding and not just the consumption of ATP through
 30 hydrolysis, is of critical importance in the helicase
 mechanism.

Although RNA unwinding activity by NPH-II is dependent on
 ATP hydrolysis (Shuman, 1992), one must establish
 35 quantitative link between ATP utilization and a specific

amount of unwinding to conclude that DExH/D proteins behave as true molecular motors (Schnapp, 1995). Plots of reaction amplitude versus ATP concentration resulted in sigmoidal curves (Fig. 12b, 12c) that are affected by duplex length: the longer the duplex, the more pronounced the sigmoidal shape of the time course. These results, together with the observation of a discrete step size and the fact that NPH-II does not dissociate in the ATP-bound state are consistent with the kinetic scheme shown in Fig. 13. This scheme was used to derive an explicit equation that describes the plots of reaction amplitudes versus ATP concentration (Fig. 12b, 12c).

$$A = (1 - x) \cdot \left(1 + \frac{k_e}{k_{u[s]}} \cdot \frac{K'_d}{[ATP]} \right)^{-Um}$$

The fit of this expression to the data establishes a direct relationship between processivity and ATP concentration, indicating that translocation is coupled to the binding of ATP. This link between ATP utilization and directional movement supports the assertion that NPH-II is a processive molecular motor.

The results suggest the following basic mechanism for RNA duplex unwinding by the DExH RNA helicase NPH-II (Fig. 13); unwinding is preceded by a slow initiation step (2.3 ± 0.2 in Co^{2+} , 3.5 ± 0.3 in Mg^{2+}). Subsequent unwinding occurs in distinct consecutive steps of roughly one-half helix turn in a defined 3'-to-5' direction with respect to the single-strand 3' overhang. The helicase

translocates rapidly along the RNA substrate, unwinding RNA with a rate that is dependent on identity of the metal ion cofactor ($13.4 \pm 0.8 \text{ min}^{-1}$ per step for Co^{2+} ; $\geq 350 \text{ min}^{-1}$ per step for Mg^{2+}). During each step, a fraction of protein, predominantly in non-ATP-bound states, dissociates from the RNA. The dependence of reaction on ATP hydrolysis and the direct connection between ATP binding and translocation indicate that NPH-II is a processive directional motor for unwinding RNA. This activity DexH/D proteins might ensure processive, regulated rearrangement of structured RNA in macromolecular assemblies during processes such as RNA splicing, export or translation initiation. A rate-limiting step before translocation would provide a straight-forward way to control unwinding by blocking and deblocking an early initiation event. Alternatively, factors that control the local concentration of ATP or accessibility of the ATP binding site on the protein could regulate processivity, enabling the helicase to dissociate from RNA at appropriate times.

DEXH/DEAD box proteins (putative RNA helicases) are important in all aspects of RNA transcription, maturation and translation. A step towards understanding the function(s) of this class of enzymes at the molecular level is to investigate the mechanism underlying duplex RNA unwinding by the vaccinia virus DEXH RNA helicase NPH-II.

Transient kinetic experiments were performed using a series of fluorescent dye labeled RNA substrates with duplex regions of different length. The unwinding reaction was monitored continuously in real time by measuring the changes in fluorescence energy transfer upon the unwinding reaction.

Timecourses under single turnover conditions displayed significant dependence on the length of the duplex region in terms of reaction rate and reaction amplitude. Reaction rates were also affected by the nature of the divalent cation cofactor. In the presence of Co^{2+} timecourses were found to have a significant lag phase, increasing with the duplex length. In the presence of Mg^{2+} no comparable lag phase was observed, but reaction rate as well as reaction amplitude were greater as compared to the reactions with Co^{2+} . Dependent on the duplex length, the ATP concentration affected the reaction rate and amplitude with both metal cofactors.

These findings suggest a basic mechanism for the translocation/unwinding reaction consisting of consecutive events of ATP binding and translocation. Quantitative description of this mechanism leading to the rate and step size for the translocation/unwinding reaction follows.

Design of the RNA substrates. These studies used the RNA helicase, vaccinia virus DExH protein NPH-II. The protein was expressed in baculavirus infected insect cells (Gross and Shuman, 1995). The NPH-II helicase requires the substrate to have single stranded overhangs 3'-end to the duplex region (Shuman, 1992), which are necessary for the helicase to bind the substrate (Shuman, 1993). In order to prevent binding of the protein at multiple sites, a substrate for mechanistical studies should contain solely one single stranded region. This single stranded region should be constant for a series of substrates where the length of the duplex region is variable.

Two different ways to design the substrates: by chemical synthesis with subsequent template directed ligation (Moore and Sharp, 1992) and by *in vitro* transcription from PCR generated templates (Chabot, 1992) with subsequent DNAzyme processing (Santoro and Joyce, 1997) in order to produce perfect blunt ended duplexes. Chemical synthesis bears the advantage of complete flexibility in the sequence choice and allows the incorporation of a variety of modifications including nucleoside and backbone modifications as well as fluorescent labels. The method is limited in terms of the length of the oligos which can be produced. *In vitro* transcription from PCR generated templates does not have these limitations for the length of the oligos but this method is not completely variable concerning the sequence and it does not allow the incorporation of extensive modifications.

Exploiting the respective advantages of both methods we designed two series of substrates. The series based on chemical synthesis contains a U-35 single stranded overhang and duplex regions from 12 to 36 basepairs (Fig. 2). The series based on *in vitro* transcription contains a 33 nucleotide overhang and duplex regions from 36 83 basepairs. The unwinding reaction of the substrates was monitored by separation of unwound strands from intact substrate on native gel electrophoresis. All substrates tested were unwound by NPH-II (Fig. 2).

Effect of trap RNA on the unwinding reaction. Excess single stranded RNA (Trap RNA) is required for single turnover reactions where dissociation of protein from the substrate has to be irreversible, i.e. trap RNA has to prevent re-binding of protein to the substrate during the

reaction (Fig. 3a). The effect of trap RNA addition with two substrates was tested: one with a 36bp duplex region and one with a 83bp duplex region. Reaction rates and amplitudes without trap addition were comparable with both substrates (Fig. 3b). The addition of trap RNA did not significantly change the reaction rates with both substrates. However, reaction amplitudes decreased with both of the substrates as compared to the reactions without trap (Fig. 3b), whereby the reaction amplitude with the longer substrate (LS83) was lower than the amplitude with the shorter substrate (LS36). Thus, the duplex length does not affect the rate of unwinding suggesting that the rate-limiting step for the overall reaction is an event which is not a potential translocation. However, the different reaction amplitudes indicate that protein dissociates off the substrate more frequently during the of unwinding of longer duplexes, suggesting that the number of dissociation "events" increases with increasing duplex length. This is an indirect indication that there are distinct consecutive steps in the unwinding reaction, even if this is not reflected in the reaction rate (Fig. 1).

Monitoring duplex unwinding by fluorescence measurements.

In order to obtain data of higher time resolution, the unwinding reaction was monitored by measuring changes in fluorescence in real time upon the unwinding of fluorescently labeled duplexes (Fig. 4a). Since the data obtained by gel shift and by fluorescence are superimposable, the real time fluorescence method provided also the control that no artifacts were introduced by the gel separation method (Fig4b). The reaction rates with both substrates tested (18bp duplex

and 36bp duplex) were similar (Fig.4b). The reaction amplitude of the 36bp duplex was lower than that of the 18bp duplex. Thus, these observations agree with the data described above (Fig. 3)

5

Effect of the ATP concentration on the unwinding reaction. The decreasing reaction amplitude with increasing duplex length in reactions where protein dissociation from the substrate was irreversible (Figs. 3 and 4) indicates that with each "unwinding step" a certain fraction of protein dissociates from the substrate (Fig. 1). For the understanding of the mechanism of unwinding it is of importance to know whether this dissociation occurs in an ATP-bound state of the protein or in a non-ATP-bound state or in both states. Experimentally, this question can be accessed by measuring the effect of ATP concentration variation on the reaction amplitude. Decreasing reaction amplitudes with decreasing ATP concentrations would indicate that dissociation occurs in a non-ATP-bound state. Dissociation in an ATP-bound state would be indicated by differences in the reaction amplitudes with duplexes of different length under ATP saturation of the system. Measurements conducted of the reaction amplitude in dependence on the ATP concentration with a 36bp substrate and a 83bp substrate (Fig. 5). The reaction amplitude decreased with decreasing ATP concentrations, indicating clearly that dissociation occurs at non-ATP bound states of the protein. The differences in the reaction amplitude of both substrates near ATP saturation suggest further that the protein dissociates also in the ATP-bound state. (Fig. 5).

10

15

20

25

30

Effect of Mg^{2+} substitution on the unwinding reaction.

At the reaction conditions tested so far, the only response to the variation in the duplex length was the reaction amplitude. Although this suggests a processive unwinding reaction, the fact that no effect was found on the rates of the unwinding reaction indicates that not a potential translocation limits the overall rate of the reaction. However, for the quantification of the unwinding steps in terms of the step size and individual rate constants it is necessary that the translocation directly affects the reaction rate. Therefore, attempts were made to decrease the rate of translocation relative to the rate limiting step such that intermediates of the unwinding reaction would be kinetically detectable and translocation would be directly reflected in the unwinding timecourse (Fig. 1). Variation in pH and temperature did not result in the desired changes in the timecourses (not shown). Subsequently, we examined whether a substitution of Mg^{2+} with other divalent cations, which act as cofactor for the ATP hydrolysis, provides reaction conditions where the translocation would be slower relative to the previously rate limiting step. Recently, only Mn^{2+} and Co^{2+} were found to substitute for Mg^{2+} in the unwinding reaction for NPH-II (Shuman, 1992). In reactions with RNA-protein pre-annealing and in the presence of trap RNA, Mn^{2+} did not change the timecourses as compared to the reactions with Mg^{2+} (Fig. 6). However, substituting Mg^{2+} with Co^{2+} resulted in a substantial change in the observed timecourse (Fig. 6). With Co^{2+} , the timecourse displayed a pronounced lag phase, suggesting reaction intermediates (Fig. 1). This assumption was supported by measuring reactions with substrates with duplex regions of different length in the presence of Co^{2+} . The timecourses

showed an increasing lag phase with increasing duplex length, i.e. the longer the duplex, the more intermediates appear in the unwinding reaction (Fig. 7). This indicated that in the presence of Co^{2+} , it can be directly observed that the reaction proceeds in distinct consecutive steps and can therefore be considered to be processive. In accordance with the results described above (Fig. 2 and 3), the reaction amplitude decreased with increasing duplex length (Fig. 7), indicating that protein dissociates more frequently during the unwinding of longer duplexes.

Employing multi-piece substrates for probing the directionality of the unwinding reaction. The results described above (Figs. 2-7) suggested that the unwinding of RNA duplexes by NPH-II is in fact processive. The second basic feature of the unwinding reaction, the directionality of the process, could not be accessed yet. Since monitoring the unwinding reaction with the described methods bears the disadvantage that the observed unwinding is caused by an event at the end of the unwinding reaction (Fig. 1), no direct evidence for directionality can be provided. Although a processive unwinding together with the binding of the protein at the single stranded overhang of the substrate indirectly argues for a certain directionality of the process, a number of alternative unwinding mechanisms (for instance looping of the single-stranded-bound protein into the duplex region) cannot be ruled out. In order to prove directionality unambiguously, we attempted to employ a more direct assay, where, beside, the unwinding of the "whole" duplex at least one intermediate has to be monitored during the reaction. We accomplished that by designing multi-piece-substrates (MPS), which comprise a

regular bottom strand but two adjacent to each other binding top strand pieces (Fig. 8a). Provided that the two pieces can be separated from each other on native PAGE, it should be possible to observe the separate displacement of the two pieces as the helicase translocates in a directional way. In order to exclude effects of the length of the different pieces *per se*, we designed two MPS; one with the longer piece next to the single stranded overhang, and another one with the shorter piece next to the single stranded overhang (Fig. 8a). The unwinding reactions were performed with protein-RNA pre-annealing and in the presence of trap RNA, such that dissociation of protein from the substrate was irreversible.

Both MPS were unwound by NPH-II, emphasizing that the helicase tolerates nicks in the top strand of the substrate. In the presence of Mg^{2+} , both pieces were displaced with a similar rate (Fig. 8b). The piece binding more distal to the overhang was displaced with a lower amplitude as compared to the piece binding closer to the single stranded overhang. This was observed for both MPS, and, therefore, the amplitude effects are caused only by the position of the pieces relative to the overhang. These results support the findings with the regular substrates (Figs. 2,3), where, although more protein dissociates from the substrate with increasing duplex length, the reaction rate is not limited by the translocation. Moreover, the fact that the amplitude was lower for the piece more distal to the overhang but the reaction rates of both pieces were similar suggests that the event which limits the rate of the overall reaction is before the translocation process, i.e. the unwinding

initiation is likely to limit the overall reaction in the presence of Mg^{2+} .

In the presence of Co^{2+} , the displacement of both pieces occurred with different lag phases (Fig. 8c). The lag for the pieces next to the overhang was always smaller, than the lag for the pieces more distal to the overhang (Fig. 8c), indicating that one piece is displaced after the other and that this displacement process has a directionality: initiating from the single stranded region the helicase translocates towards the other end of the duplex while it unwinds it. Moreover, the reaction amplitude for the piece more distal to the overhang was smaller than the amplitude for the piece next to the overhang, indicating that also in the presence of Co^{2+} protein dissociates from the substrate more frequently with increasing duplex length.

Materials and Methods

NPH-II was expressed and purified as described (Gross and Shuman, 1995). RNAs were prepared by *in vitro* transcription of PCR-generated T7 transcription templates (Jankowsky and Schwenzer, 1996) or by chemical synthesis (Wincott et al., 1995). The PCR-amplified DNA templates represent a segment of the ampicillin-resistance gene of the pBS(+/-) phagemid (Stratagene). To generate perfect blunt-end duplexes, the RNA pieces were trimmed by site-directed processing using engineered DNazymes (Santoro and Joyce, 1997; Pyle et al., 2000) with 12 nucleotides on each of the binding arms. Bottom-strand RNAs were joined from two separately synthesized oligonucleotides by template-directed ligation using T4 DNA ligase (Moore and Sharp, 1992) RNA duplexes were formed by combining the bottom-strand RNA with a five-fold molar excess of γ^{32}

P-labelled top strand in 10mM MOPS, 6.5, 1mM EDTA. The solution was heated to 95°C for 2 min and cooled to room temperature over 90 min. Duplexes were separated from single-strand RNA by native PAGE, visualized by radiolytic scanning (Packard Instant Imager) and excised from the gel.

Unwinding reactions

Unwinding reactions were performed at room temperature in 30µl of 40mM Tris-HCl buffer, pH 8.0 and 4mM MgCl₂ or CoCl₂. The reaction also contained 25mM NaCl, which was introduced with the protein storage buffer. In a typical reaction, 1-2nM RNA substrate was incubated with 10-15nM NPH-II in reaction buffer without ATP at room temperature for 5-7 min. Longer pre-incubation time did not change the reaction kinetics. Saturation of substrate with protein before the reaction was verified by gel-shift analysis (Lohmann and Bjoernson, 1996). The unwinding reaction initiated by adding the ATP to a final concentration of 3.5mM unless otherwise stated. Aliquots at the respective time points were quenched with two volumes of stop buffer 25 mM EDTA, 0.4% SDS, 0.05% BPB, 0.05% XCB, 10% glycerol containing 200nM of unlabelled top-strand RNA to prevent re-annealing of unwound duplexes during electrophoresis. Reannealing of unwound duplexes during the reaction did not affect unwinding kinetics under any reaction conditions as verified by independent measurements of association rate constants of the substrate strands. Excess RNA (Trap RNA) (200 nM final concentration added with the ATP) was a single-strand of 39-nucleotides of unrelated sequence. Higher concentrations of trap RNA did not change observed reaction amplitudes.

Unwinding reactions were also monitored in real time using fluorescence energy transfer experiments on substrates labelled at the 3' and 5' ends of the duplex terminus farthest from the single stranded 3' overhang. These measurements were superimposable with results from gel-shift measurements and confirmed the first-order nature of the time courses as well as the rate constants.

Kinetic analysis

Data were fit numerically using the FITSIM software package (Barshop and Frieden, 1983). Time courses were corrected for the reaction amplitudes, and initial kinetic parameters were determined using the KINSIM software (Barshop and Frieden, 1983). Unwinding step size and rate constants were determined from time courses in Co^{2+} using a kinetic model in which the unwinding reaction consists of N consecutive first-order reactions where N is the number of kinetic steps. Rate constants (except for the first step see below) were linked, that is, forced to yield the same value in the fitting. The linked rate constants and the first rate constant were allowed to float simultaneously. On the basis of the observation that an initial step is slower than subsequent steps in Mg^{2+} (see Fig. 11b and text), the first kinetic step was defined as having a different rate constant (k_1) than subsequent kinetic steps (k_u). Using this model, increasing numbers of steps (N) were examined iteratively for each duplex until k_1 and k_u approached constant values for the different duplexes. Convergence occurred when $k_1 = 2.3 \pm 0.2 \text{ min}^{-1}$ and $k_u = 13.4 \pm 0.8 \text{ min}^{-1}$.

Note that k_u is an apparent rate constant that depends on the ATP concentration according to $k_u = k_{u[S]} [\text{ATP}/(\text{ATP} + K'_d)]$, where $k_{u[S]}$ is the respective rate constant at ATP

saturation. The step size (m) was calculated according to $m=L(N-y)$, where y is the number of kinetic steps do not contribute to the actual strand displacement and L is the duplex length. A step size of $m=6$ and a value of $y=1$ were consistently calculated for all duplexes λ including multi-piece substrates.

The lower limit for the translocation rate constant in Mg^{2+} was estimated by considering that a lag phase was not observed with the longest (83 bp) duplex (Fig. 9b). Assuming an unwinding step size identical to that in Co^{2+} (6 bp), and that k_u must be equal or greater than a value that does not cause an apparent lag phase in 14 unwinding steps, the translocation rate constant was estimated as $k_u > \sim 350 \text{ min}^{-1}$.

Equation (1) was derived from the expression describing the dependence of reaction amplitude (A) on the processivity (P):

$$A = (1 - x) P^{1/m}$$

where x is the fraction of protein that dissociates from the substrate before unwinding occurs. P is then substituted with a term that relates processivity and ATP concentration under the conditions specified by the scheme in Fig. 13.

5

$$p = \frac{k_{\text{on}} \left(\frac{[\text{ATP}]}{K'_d + [\text{ATP}]} \right)}{k_{\text{on}} \left(\frac{[\text{ATP}]}{K'_d + [\text{ATP}]} \right) + k_r \left(1 - \frac{[\text{ATP}]}{K'_d + [\text{ATP}]} \right)}$$

10

where K'_d is the apparent dissociation constant of ATP from the enzyme substrate complex.

RNA Substrate Preparation

15

Chemical Synthesis: RNA oligonucleotides were prepared by chemical synthesis using phosphoramidite chemistry (amidites purchased from Glen Research) on an ABI 392 RNA/DNA synthesizer. Deprotection of the crude oligonucleotides was carried out according to standard protocols (Wincott et al., 1995). All RNA oligonucleotides were purified on denaturing PAGE.

20

25

The bottom strand RNAs were joined out of two separately synthesized oligonucleotides by template directed ligation using T4 DNA ligase (Moore and Sharp, 1992). The ligated products were purified on denaturing PAGE.

30

Fluorescent Labeling: Amino linkers were placed in the 3' or 5' position of the respective RNA during the synthesis. After removal of all deprotection groups, the modified RNA was labeled with fluorescein-or rhodamine-isothiocyanate (Sigma), respectively. Labeling reactions were carried out in the dark for 12-18 hours with 50 mM

dye-isothiocyanate and RNA concentrations below 50 μ M in 0.1 M sodiumbicarbonate (pH 9.0) and 20% (v/v) DMSO for the fluorescein-isothiocyanate labeling and 50% (v/v) DMSO for the rhodamine-isothiocyanate labeling. The excess rhodamine-isothiocyanate was removed by repeated phenol extractions followed by ethanol precipitation. The excess fluorescein-isothiocyanate was removed by repeated ethanol precipitations. All labeled RNAs were finally purified on denaturing PAGE.

In vitro transcription: RNAs for the multi-piece-substrates were prepared by *in vitro* transcription of PCR generated templates (Chabot, 1992). The PCR-amplified DNA-templates represent a segment of the ampicillin resistance gene of the pBS (+/-) phageimid (Stratagene). Templates for bottom and top strand RNAs were separately amplified from ScaI linearized phageimid using different pairs of primers with the respective sense primer containing the binding sequence as well as the T7 promoter sequence. 100 μ L PCR reactions were carried out for 30 cycles (52 $^{\circ}$ C, 72 $^{\circ}$ C, 94 $^{\circ}$ C, 1 min each) followed by a 7 min final extension at 72 $^{\circ}$ C. Homogeneity of the PCR products was verified on a 2% agarose gel. The PCR reaction solutions were extracted with an equal volume of phenol, followed by extraction with chloroform isoamylalcohol (Sambrook et al., 1989). Not incorporated dNTPs were removed by SEC (NAP-5, Pharmacia). Subsequently, the volume of the solution was reduced in a Speedvac concentrator (Savant). The obtained DNA templates were used for *in vitro* transcriptions with T7 polymerase according to standard procedures. The transcribed RNAs were purified on denaturing PAGE.

The top strands for the Multi-Piece-Substrates were produced by site directed processing using an engineered DNAzyme (Santoro and Joyce, 1997). DNAzymes with 12 nucleotides in each of the binding arms were designed to cut at the desired positions. The DNAzyme reactions (10 μ M RNA, 30 μ M DNAzyme) were carried out for 4h at 40 $^{\circ}$ C followed by ethanol precipitation. Subsequently, the DNAzyme was destroyed by RQ RNase free DNase (Promega) in a 100 μ L volume with 15units DNase for 60 min at 37 $^{\circ}$ C. The RNA was ethanol precipitated and the products were purified by denaturing PAGE.

Duplexes were formed by combining the bottom strand RNA with a 5 fold molar excess of labeled (either radioactively or fluorescently) top strand in 10mM MOPS (pH 6.5), 1mM EDTA. The solution was heated to 95 $^{\circ}$ C and cooled to room temperature within 90 min. Duplexes were separated from single stranded RNA on native PAGE. The RNA was visualized either by UV-shadowing or by radiolytic scanning (Packard Instant Imager), cut out, eluted, and ethanol precipitated.

Helicase reactions

Unwinding reactions were carried out at room temperature in 40 mM Tris/HCl (pH 8.0) and 3mM MgCl₂ (MnCl₂, CoCl₂), 20mM NaCl was present in the reaction due to introduction with the protein storage buffer. In a typical reaction, 3nM RNA substrate was incubated with 10-15 nM NPH-II in reaction buffer without ATP at room temperature for 7 min. Longer incubation time did not change the observed reaction kinetics. The reaction was started by adding the ATP. In reactions with trap RNA, the trap was added together with the ATP at the reaction start.

Gel shift measurements: For the reactions monitored by gel shift PAGE, reactions were performed in a volume of 30 μ L. Aliquotes were taken at appropriate times and combined with two volume of stop buffer (25mM EDTA, 0.4% SDS, 0.05% BPB, 0.05% XCB, 10% glycerol), containing 200nM of unlabeled top strand in order to prevent re-annealing of unwound duplexes during electrophoresis. Unwound strands were separated from intact substrate on native PAGE and the amount of unwound duplex was determined using a Molecular Dynamics PhosphorImager and the Imagequant Software.

Fluorescence Measurements: For the reactions monitored by fluorescence measurements, reactions were performed in fluorescence cuvette (600 μ L) in an Aminco SLM AB2 fluorescence spectrometer. Excitation wavelength was set at 492 nm, (8nm slid width) emission was recorded at 518 nm (8 nm slid width, PMT sensitivity 600-800V). The reaction was started as described and measurements were taken automatically in fixed intervals.

Employing a series of designed model substrates, it was shown thar the unwinding of RNA duplexes by the DExH protein NPH-II is processive. In the presence of Mg^{2+} the translocation does not limit the overall reaction rate. The rate-limiting step of the unwinding reaction is likely to be the unwinding initiation. The sensitivity of the reaction amplitude to the duplex length indicates that during each "unwinding step" a certain fraction protein dissociates from the substrate. The dependence of the reaction amplitude on the ATP concentration emphasizes that this dissociation occurs in non-ATP bound states of the protein as well as in ATP-bound states of the protein.

Also developed was an assay to test the directionality of the unwinding process and showed that the unwinding process has a defined directionality: initiating from the single stranded region the helicase translocates towards the other end of the duplex while it unwinds it. The proof of the processivity and directionality of the unwinding process is the prerequisite for quantitative characterization of the helicase reaction which is necessary to derive functional models.

5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

References

Ali, J.A., and Lohmann, T.M. (1997) "Kinetic Measurement of the Step Size DNA Unwinding by *Escherischia coli* UvrD Helicase", *Science*, 275, 377-380.

5

Barshop, B. A. and Frieden, C. (1983) "Analysis of Numerical Methods For Computer Simulation of Kinetic Processes: Development of KINSIM--A Flexible Portable System. *Anal. Biochem.* 130, 134-145.

10

Chabot, B. (1992) "Synthesis and Purification of RNA Substrates, in *RNA Processing: A Practical Approach*", Higgins, S.J. and Hames, B.D. Eds., IRL Press, Oxford, U.K., pp 1-31.

15

de la Cruz, I., Kresler, D. and Linder P. (1999) "Unwinding RNA in *Saccharomyces Cerevisiae*: DEAD-Box Proteins and Related Families," *Trends Biochem. Sc.* 24, 192-198.

20

Gross, C.H. and Shuman, S. (1995) "Mutational Analysis of Vaccinia Virus Nucleoside Triphosphate Hydrolase II, a DExH Box RNA Helicase", *J. Virol.*, 69, 4727-4736.

25

Gross, C.H. and Shuman, S. (1996) "Vaccinia Virus RNA Helicase: Nucleic Acid Specificity in Duplex Unwinding, *J. Virol*, 70, 2615-2619.

30

Jankowsky, E. and Schwenzer, B. (1996) "Efficient Improvement of Hammerhead Ribozyme Mediated Cleavage of Long Substrates by Oligonucleotide Facilitators" *Biochemistry* 37, 15313-15321.

Jeltsch, A. and Pingoud, A. (1998) "Kinetic Characterization of Linear Diffusion of the Restriction Endonuclease EcoRV on DNA, Biochemistry 24, 2160-2169.

5 Kadare, G. and Haenni, A. L., (1997) "Virus encoded RNA Helicases," J. Virol 71, 2583-2590.

10 Kelemen, B. R. and Raines, R. T. (1999) "Extending the Limits to Enzymatic Catalysis: Diffusion of Ribonuclease A in One Dimension" Biochemistry 27, 5302-5307.

15 Lohmann, T. M. and Bjoernson, K.P. (1996) "Mechanism of Helicase Catalyzed DNA Unwinding", Annu. Rev. Biochem. 65, 169-214.

20 Moore, M.J. and Sharp, P.A. (1992) "Site Specific Modification of Pre-Messenger RNA-the 2'-Hydroxyl Groups at the Splice Sites", Science, 256, 992-997.

25 Pyle, A. M., et al., (2000) "Using DNazymes to Cut, Process and Map RNA Molecules For Structural Studies or Modification" Methods Enzymol 317, 140-146.

30 Santoro, S.W. and Joyce, G.F. (1997) "a General Purpose RNA-Cleaving DNA Enzyme", Proc. Natl. Acad. Sci., USA, 94, 4262-4266.

Schnapp, B. Molecular Motors, (1995) "Two Heads Are Better Than one" Nature 373, 655-666.

35 Shimamoto, N. (1999) "One-Dimensional Diffusion of Proteins Along DNA, J. Biol. Chem. 274, 15239-15296.

Shuman, S. (1992) "Vaccinia Virus RNA Helicase: An Essential Enzyme Related to the DE-H Family of RNA Dependent NTPases", Proc. Natl. Acad. Sci. USA, 89, 10935-10939.

5

Shuman, S. (1993) "Vaccinia Virus RNA Helicase, J. Biol. Chem., 268, 11798-11802.

10

Staley, J. P. and Guthrie, C. (1998) "Mechanical Devices of the Spliceosome: Motors Clocks Springs and Things" Cell 90, 1041-1050.

15

Wagner, I.D.O., et al., (1998) "The DEAH-Box Protein Prp 22 is an ATPase that Mediates ATP Dependant mRNA Release From the Spliceosome and Unwinds RNA Duplexes," EMBO J. 17, 2926-2937.

20

Wincott et al. (1995) "Synthesis, Deprotection, Analysis and Purification of RNA and Ribozymes", Nucleic Acids Res., 23, 2677-2684.

25

Xiang, Q., et al., (1998) "Sequence Specificity of a Group II Intron Ribozyme: Multiple Mechanisms For Promoting Unusually High Discrimination Against Mismatched Targets" Biochemistry 37, 3839-3849.

30

Zimmeric, C. and Frieden, C. (1998) "Analysis of Progress Curves by Simulations Generated by Numerical Integration" Biochem. I. 258, 381-387.